

## SCIENTIFIC NOTE

**MCOL, frontalin and ethanol: A potential operational trap lure for Douglas-fir beetle in British Columbia****B. Staffan Lingren<sup>1,2</sup>, Daniel R. Miller<sup>3</sup>, J. P. LaFontaine<sup>4</sup>**

The Douglas-fir beetle, *Dendroctonus pseudotsugae* (Coleoptera: Curculionidae) is a major pest of Douglas-fir, *Pseudotsuga menziesii* (Mirb.) in British Columbia (Humphreys 1995). An operational trap lure for *D. pseudotsugae* could be useful in an integrated pest management program to minimize mortality of Douglas-fir, particularly in conjunction with anti-aggregation pheromones (Lindgren *et al.* 1988; Ross and Daterman 1995a). The principal pheromone of *D. pseudotsugae* is frontalin (1,5-dimethyl-6,8-dioxabicyclo [3.2.1] octane), which is produced by male and female beetles and attracts both sexes of beetles (Pitman and Vité 1970; Kinzer *et al.* 1971; Rudinsky *et al.* 1976). In British Columbia, *D. pseudotsugae* prefer multiple-funnel traps baited with racemic frontalin (50:50 mix of the two enantiomers) or (S)-(-)-frontalin equally over those baited with (R)-(+)-frontalin (Lindgren 1992).

Two additional pheromones are produced by female *D. pseudotsugae* and attract both sexes of beetles, particularly when presented with host odours or frontalin: MCOL (1-methylcyclohex-2-en-1-ol) (Libbey *et al.* 1983; Lindgren *et al.* 1992; Ross and Daterman 1995b) and seudenol (3-methylcyclohex-2-en-1-ol) (Vité *et al.* 1972; Rudinsky *et al.* 1974; Pitman *et al.* 1975; Ross and Daterman 1995b). These two compounds are isomers of each other.

In 1991, we conducted a trapping experiment in British Columbia, targeting *D. pseudotsugae*. The objective of the experiment was to determine the effect of racemic frontalin and racemic MCOL, alone and in combination, on the attraction of *D. pseudotsugae* to traps baited with ethanol in

British Columbia. Lindgren *et al.* (1992) found that the attraction of beetles to traps baited with the two enantiomers of MCOL seem to be additive with the highest catches in traps baited with racemic MCOL. In laboratory assays, ethanol enhanced the activity of frontalin on arrestment of male *D. pseudotsugae* (Libbey *et al.* 1983). In field assays, ethanol increased catches of beetles in traps baited with frontalin and seudenol (Pitman *et al.* 1975; Ross and Daterman 1995b).

PheroTech Inc. (now Contech, Victoria, BC) supplied all traps and lures. Chemical purities were >95% for all semiochemicals. Release rates were determined gravimetrically at 20–23 °C. Traps were suspended from a metal pole made from electrical conduit tubing such that the bottom of each trap was 0.2–0.5 m above ground level. No trap was suspended within 2 m of any tree. All lures were placed within the funnels (Lindgren 1983).

The experiment was conducted in mature stands of Douglas-fir at three locations in southern British Columbia: 1) Maple Ridge (10 April–12 May 1991); 2) Cache Creek (13–31 May 1991); and 3) Invermere (30 May–8 August 1991). We used 40 12-unit multiple-funnel traps (Lindgren 1983) with dry cups in Maple Ridge and Cache Creek, whereas 20 traps were used in Invermere. Each collecting cup contained a small piece of Vapona No-Pest Strip (Green Cross; Fisons Horticulture, Mississauga, Ontario, Canada) as a killing agent to prevent damage to the target species by predatory insects. At each location, traps were set in blocks of four traps per block resulting in 10, 10, and 5 replicate blocks per location, respectively. Blocks, and traps within

<sup>1</sup> Corresponding author, Staffan.Lindgren@unbc.ca

<sup>2</sup> Natural Resources and Environmental Studies Institute, University of Northern British Columbia, 3333 University Way, Prince George, British Columbia, Canada V2N 4Z9

<sup>3</sup> USDA Forest Service, Southern Research Station, 320 Green Street, Athens, GA, USA 30602-2044


<sup>4</sup> Contech Enterprises Inc., 7572 Progress Way, Delta, British Columbia, Canada V4G 1E9

blocks, were spaced 10–15 m apart in Maple Ridge and Cache Creek and 50m apart in Invermere. Each trap was baited with a white PVC sleeve pouch (40 cm) releasing ethanol at approximately 53 mg/d. Racemic frontalin and racemic MCOL were released from micro-centrifuge tubes (250 µL) and plastic bubblecaps, respectively, each at a rate of approximately 2–3 mg/d. One of the following four treatments was randomly assigned to each trap within a block: (1) untreated control; (2) MCOL; (3) frontalin; and (4) MCOL + frontalin.

Data were analyzed with the SYSTAT (ver. 11) statistical package (SYSTAT Software Inc., Point Richmond, California). Trap catch data were transformed by  $\ln(y+1)$  to reduce heteroscedasticity. Data at each location were subjected to ANOVA using the following model: (1) replicate; (2) MCOL; (3) frontalin; and (4) MCOL x frontalin.

Trap catches of *D. pseudotsugae* were significantly affected by MCOL and frontalin at all three locations (Fig. 1). The responses were additive in Maple Ridge and Cache Creek, as there was no significant interaction with MCOL and frontalin at either location (Fig. 1A–B). There was a significant MCOL x frontalin interaction on trap catches in Invermere, resulting in a synergistic effect (Fig. 1C). The difference may be due to the relatively close spacing of treatments in the Maple Ridge and Cache Creek experiments. In contrast to our results, Ross and Daterman (1995b) found that MCOL had no effect on catches of *D. pseudotsugae* in traps baited with ethanol and

frontalin in Oregon. In their study, the combination of ethanol, frontalin, and seudenol was the most effective lure combination for Douglas-fir beetles. Geographic variation in chemical ecology is known for *D. pseudotsugae*, warranting further trials of trap-lure blends over a broader range (Ryker *et al.* 1979; Stock *et al.* 1979; Ross and Daterman 1995b). Nevertheless, traps baited with frontalin, MCOL, and ethanol as described here should be used for trapping



**Figure 1.** Effect of MCOL and frontalin on catches of *D. pseudotsugae* in traps baited with ethanol in Maple Ridge (A), Cache Creek (B), and Invermere (C). Significance levels (P) for ANOVA on trap catches.

Douglas-fir beetles in British Columbia.

ACKNOWLEDGEMENTS

We thank Emile Begin for field work in Invermere; the University of British Columbia for permission to conduct the study in the Maple Ridge Research Forest; and Dave Piggitt for help locating sites near Cache Creek. Funding for this research was provided by NSERC (BSL), a grant (BSL) and an Industrial Post-doctoral Fellowship (DRM) from the Science Council of British Columbia,

and from the USDA Forest Service (DRM). The use of trade names and identification of firms or corporations does not constitute an official endorsement or approval by the United States Government of any product or service to the exclusion of others that may be suitable. The USDA is an equal opportunity provider and employer.

## REFERENCES

- Humphreys, N. 1995. Douglas-fir beetle in British Columbia. Natural Resources Canada, Canadian Forest Service Forest Pest Leaflet 14.
- Kinzer, G. W., A. F. Fentiman, Jr., R. L. Foltz, and J. A. Rudinsky. 1971. Bark beetle attractants: 3-Methyl-2-cyclohexen-1-one isolated from *Dendroctonus pseudotsugae*. Journal of Economic Entomology 64: 970–971.
- Libbey, L. M., A. C. Oehlschlager, and L. C. Ryker. 1983. 1-Methylcyclohex-2-en-1-ol as an aggregation pheromone of *Dendroctonus pseudotsugae*. Journal of Chemical Ecology 9: 1533–1541.
- Lindgren, B. S. 1983. A multiple funnel trap for scolytid beetles (Coleoptera). The Canadian Entomologist 115: 299–302.
- Lindgren, B. S. 1992. Attraction of Douglas-fir beetle, spruce beetle and a bark beetle predator (Coleoptera: Scolytidae and Cleridae) to enantiomers of frontalin. Journal of the Entomological Society of British Columbia 89: 13–17.
- Lindgren, B. S., M. D. McGregor, R. D. Oakes, and H. E. Meyer. 1988. Effect of MCH and baited Lindgren traps on Douglas-fir beetle attacks on felled trees. Journal of Applied Entomology 105: 289–294.
- Lindgren, B. S., G. Gries, H. D. Pierce, Jr., and K. Mori. 1992. *Dendroctonus pseudotsugae* Hopkins (Coleoptera: Scolytidae): Production and response to enantiomers of 1-methylcyclohex-2-en-ol. Journal of Chemical Ecology 18: 1201–1208.
- Pitman, G. B., and J. P. Vité. 1970. Field response of *Dendroctonus pseudotsugae* (Coleoptera: Scolytidae) to synthetic frontalin. Annals of the Entomological Society of America 63: 661–664.
- Pitman, G. B., R. L. Hedden, and R. I. Gara. 1975. Synergistic effects of ethyl alcohol on the aggregation of *Dendroctonus pseudotsugae* (Col., Scolytidae) in response to pheromones. Zeitschrift für angewandte Entomologie 78: 203–208.
- Ross, D. W., and G. E. Daterman. 1995a. Pheromone-based strategies for managing the Douglas-fir beetle on a landscape scale, pp. 347–358. In F. P. Hain, S. M. Salom, W. F. Ravlin, T. L. Payne, and K. F. Raffa (eds.), Proceedings: Behavior, population dynamics and control of forest insects, Joint IUFRO Working Part Conference, Maui, Hawaii, February 6–11 1994. Ohio State University, Wooster, OH.
- Ross, D. W., and G. E. Daterman. 1995b. Response of *Dendroctonus pseudotsugae* (Coleoptera: Scolytidae) and *Thanasimus undatulus* (Coleoptera: Cleridae) to traps with different semiochemicals. Journal of Economic Entomology 88: 106–111.
- Rudinsky, J. A., M. E. Morgan, L. M. Libbey, and T. B. Putnam. 1974. Additional components of the Douglas fir beetle (Col., Scolytidae) aggregative pheromone and their possible utility in pest control. Zeitschrift für angewandte Entomologie 76: 65–77.
- Rudinsky, J. A., M. E. Morgan, L. M. Libbey, and T. B. Putnam. 1976. Release of frontalin by male Douglas-fir beetle. Zeitschrift für angewandte Entomologie 81: 267–269.
- Ryker, L. C., L. M. Libbey, and J. A. Rudinsky. 1979. Comparison of volatile compounds and stridulation emitted by the Douglas-fir beetle from Idaho and western Oregon populations. Environmental Entomology 8: 789–798.
- Stock, M. W., G. B. Pitman, and J. D. Guenther. 1979. Genetic differences between Douglas-fir beetles (*Dendroctonus pseudotsugae*) from Idaho and coastal Oregon. Annals of the Entomological Society of America 72: 394–397.
- Vité, J. P., G. B. Pitman, A. F. Fentiman, Jr., and G. W. Kinzer. 1972. 3-Methyl-2-cyclohexen-1-ol isolated from *Dendroctonus*. Naturwissenschaften 58: 469.